

Theoretical issues with ΔS in three-body B decays

Hai-Yang Cheng*

*Institute of Physics, Academia Sinica,
Taipei, Taiwan, 115, R.O.C.*

Theoretical uncertainties with the time-dependent CP asymmetries ΔS and A_{CP} in $B^0 \rightarrow K^+K^-K_S$, $K_SK_SK_S$ and $K_S\pi^0\pi^0$ decays are discussed. In order to have a reliable estimate of CP asymmetries, it is very crucial to understand the underlying mechanism for the nonresonant background which dominates the KKK modes.

I. INTRODUCTION

Time-dependent mixing-induced and direct CP asymmetries S_f and A_f , respectively, in charmless three-body decays of the B mesons have been measured by BaBar and Belle for $K_SK_SK_S$ and $K^+K^-K^0$ final states. In addition, BaBar has also measured time-dependent CP asymmetries for $K^+K^-K_L$ and $K_S\pi^0\pi^0$ and performed a Dalitz plot analysis for $K^+K^-K^0$.

For CP eigenstates, the measured mixing-induced CP asymmetry S_f can be used to define an effective $\sin 2\beta$ via

$$S_f \equiv -\eta_f \sin 2\beta_{\text{eff}} \quad (1)$$

with η_f being the CP eigenvalue of the final state f . While $K_SK_SK_S$, $K_SK_SK_L$ and $K_S\pi^0\pi^0$ have fixed CP -parities with $\eta_f = 1, -1, 1$ respectively, $K^+K^-K_S$ is an admixture of CP -even and CP -odd components, rendering its CP analysis more complicated. By excluding the major CP -odd contribution from ϕK_S , the 3-body $K^+K^-K_S$ final state is primarily CP -even. A measurement of the CP -even fraction f_+ in the $B^0 \rightarrow K^+K^-K_S$ decay yields $f_+ = 0.89 \pm 0.08 \pm 0.06$ by BaBar [1] and $0.93 \pm 0.09 \pm 0.05$ by Belle [2], while the CP -odd fraction in $K^+K^-K_L$ is measured to be $f_- = 0.92 \pm 0.33^{+0.13}_{-0.14} \pm 0.10$ by BaBar [3]. Hence, while $\eta_f = 1$ for the $K_SK_SK_S$ mode, $\eta_f = 2f_+ - 1$ for $K^+K^-K_S$ and $\eta_f = -(2f_- - 1)$ for $K^+K^-K_L$.

The world average of $\sin 2\beta$ measured in the decays $B \rightarrow \phi_{c\bar{c}}K_S$ with $\phi_{c\bar{c}}$ being a charmonium is $\sin 2\beta_{\phi_{c\bar{c}}K_S} = 0.675 \pm 0.026$ [4]. However, the time-dependent CP asymmetries in the $b \rightarrow s\bar{q}\bar{q}$ penguin-induced two-body decays such as $B^0 \rightarrow (\phi, \omega, \pi^0, \eta', f_0)K_S$ measured by BaBar and Belle indicate $\sin 2\beta_{\text{eff}} = 0.53 \pm 0.05$ [4]. This seems to imply a 2.6σ deviation from the expectation of the standard model (SM) where CP asymmetry in all above-mentioned modes should be equal to $\sin 2\beta_{\phi_{c\bar{c}}K_S}$ [4] with a small deviation at most $\mathcal{O}(0.1)$ [5].

The measured results of $\sin 2\beta_{\text{eff}}$ for $K^+K^-K_{S,L}$, $K_SK_SK_S$ and $K_S\pi^0\pi^0$ are listed in Table IV below. In order to see if the current measurements of the deviation

of $\sin 2\beta_{\text{eff}}$ in these modes from $\sin 2\beta_{\phi_{c\bar{c}}K_S}$ signal New Physics in $b \rightarrow s$ penguin-induced modes, it is of great importance to examine and estimate how much of the deviation of $\sin 2\beta_{\text{eff}}$ is allowed in the SM. One of the major uncertainties in the dynamic calculations lies in the hadronic matrix elements which are nonperturbative in nature. One way to circumvent this difficulty is to impose SU(3) flavor symmetry [6, 7] or isospin and U-spin symmetries [8] to constrain the relevant hadronic matrix elements. While this approach is model independent in the symmetry limit, deviations from that limit can only be computed in a model dependent fashion. In addition, it may have some weakness as discussed in [7].

As mentioned above, BaBar has performed a Dalitz plot analysis for $B^0 \rightarrow K^+K^-K^0$ decays and found that it is dominated by S -wave nonresonant contributions with fraction $91 \pm 19\%$ [9]. A large nonresonant component is also found in $B^+ \rightarrow K^+K^+K^-$ decays [10, 11]. This is a surprise because it is known that in three-body decays of D mesons, the nonresonant background is at most 10%. Hence, it is very important to take into account the large nonresonant effects in any realistic model calculations.

II. FORMULISM FOR CHARMLESS 3-BODY B DECAYS

We consider the decay $\bar{B}^0 \rightarrow K^+K^-\bar{K}^0$ as an illustration. Under the factorization approach, its decay amplitude consists of three distinct factorizable terms: (i) the meson emission process, $\langle \bar{B}^0 \rightarrow K^+\bar{K}^0 \rangle \times \langle 0 \rightarrow K^- \rangle$, (ii) the transition process, $\langle \bar{B}^0 \rightarrow \bar{K}^0 \rangle \times \langle 0 \rightarrow K^+K^- \rangle$, and (iii) the annihilation process $\langle \bar{B}^0 \rightarrow 0 \rangle \times \langle 0 \rightarrow K^+K^-\bar{K}^0 \rangle$, where $\langle A \rightarrow B \rangle$ denotes a $A \rightarrow B$ transition matrix element.

For the kaon emission process, the amplitude induced by $V-A$ currents reads

$$\begin{aligned} A_{\text{current-ind}} &\equiv \langle K^-(p_3) | (\bar{s}u)_{V-A} | 0 \rangle \\ &\times \langle \bar{K}^0(p_1) K^+(p_2) | (\bar{u}b)_{V-A} | \bar{B}^0 \rangle \\ &= -\frac{f_K}{2} [2m_3^2 r + (m_B^2 - s_{12} - m_3^2) \omega_+ \\ &+ (s_{23} - s_{13} - m_2^2 + m_1^2) \omega_-], \end{aligned} \quad (2)$$

*Electronic address: phcheng@phys.sinica.edu.tw




FIG. 1: Point-like and pole diagrams responsible for the $\overline{B}^0 \rightarrow K^+ K^- K^0$ matrix element of the current $\bar{u}\gamma_\mu(1-\gamma_5)b$, where the symbol $\bullet$ denotes an insertion of the current.

where $s_{ij} \equiv (p_i + p_j)^2$, $(\bar{q}_1 q_2)_{V-A} \equiv \bar{q}_1 \gamma_\mu (1 - \gamma_5) q_2$, and r , ω_+ and ω_- are three unknown form factors. In principle, one can apply heavy meson chiral perturbation theory (HMChPT) to evaluate the form factors r , ω_+ and ω_- [12]. The relevant diagrams are depicted in Fig. 1. However, this will lead to too large nonresonant decay rates in disagreement with experiment [13]. A direct calculation indicates that the branching ratio of $B^0 \rightarrow K^+ K^- K^0$ arising from the current-induced process alone is already at the level of 77×10^{-6} which substantially exceeds the measured total branching ratio of 25×10^{-6} [4]. The issue has to do with the applicability of HMChPT. In order to apply this approach, two of the final-state pseudoscalars have to be soft. The momentum of the soft pseudoscalar should be smaller than the chiral symmetry breaking scale of order 1 GeV. For 3-body charmless B decays, the available phase space where chiral perturbation theory is applicable is only a small fraction of the whole Dalitz plot. Therefore, it is not justified to apply chiral and heavy quark symmetries to a certain kinematic region and then generalize it to the region beyond its validity.

To circumvent the aforementioned difficulty, we assume the momentum dependence of the nonresonant amplitude in the exponential form

$$A_{\text{current-ind}} = A_{\text{current-ind}}^{\text{HMChPT}} e^{-\alpha_{\text{NR}} p_B \cdot (p_1 + p_2)} e^{i\phi_{12}}, \quad (3)$$

so that the HMChPT results are recovered in the soft meson limit $p_1, p_2 \rightarrow 0$. That is, the nonresonant amplitude in the soft meson region is described by HMChPT, but its energy dependence beyond the chiral limit is governed by the exponential term $e^{-\alpha_{\text{NR}} p_B \cdot (p_1 + p_2)}$. The parameter α_{NR} can be constrained by the decay $B^- \rightarrow \pi^+ \pi^- \pi^-$ [14].

For the kaon emission process, there exist two more amplitudes, one induced by currents and the other induced by scalar densities:

$$\begin{aligned} A_1 &= \langle \overline{K}^0(p_3) | (\bar{s}d)_{V-A} | 0 \rangle \langle K^+(p_2) K^-(p_3) | (\bar{d}b)_{V-A} | \overline{B}^0 \rangle, \\ A_2 &= \langle \overline{K}^0(p_3) | \bar{s}d | 0 \rangle \langle K^+(p_2) K^-(p_3) | \bar{d}b | \overline{B}^0 \rangle. \end{aligned} \quad (4)$$

Although the 3-body matrix elements in A_1 and A_2 are

OZI suppressed, they do receive intermediate vector meson and scalar pole contributions, respectively. For example,

$$\begin{aligned} &\langle K^+(p_2) K^-(p_3) | (\bar{d}b)_{V-A} | \overline{B}^0 \rangle^R = \\ &\sum_i \frac{g^{f_{0i} \rightarrow K^+ K^-}}{m_{f_{0i}}^2 - s_{23} - im_{f_{0i}}} \langle f_{0i} | (\bar{d}b)_{V-A} | \overline{B}^0 \rangle, \end{aligned} \quad (5)$$

where f_{0i} denote the generic f_0 -type scalar mesons, $f_{0i} = f_0(980), f_0(1370), f_0(1500), X_0(1550), \dots$.

For the transition amplitudes, we need to consider the two-body matrix elements $\langle K^+ K^- | V_\mu | 0 \rangle$ and $\langle K^+ K^- | \bar{s}s | 0 \rangle$. Both receive resonant and nonresonant contributions. The first matrix element can be related to the kaon electromagnetic (e.m.) form factors $F_{em}^{K^+ K^-}$ and $F_{em}^{K^0 \overline{K}^0}$ for the charged and neutral kaons, respectively. The nonresonant form factor can be constrained by the asymptotic behavior implied by pQCD [15]. The second matrix element has the expression

$$\begin{aligned} \langle K^+(p_2) K^-(p_3) | \bar{s}s | 0 \rangle &= \sum_i \frac{m_{f_{0i}} \bar{f}_{f_{0i}}^s g^{f_{0i} \rightarrow K^+ K^-}}{m_{f_{0i}}^2 - s_{23} - im_{f_{0i}} \Gamma_{f_{0i}}} \\ &+ f_s^{NR}, \end{aligned} \quad (6)$$

where the scalar decay constant $\bar{f}_{f_{0i}}^s$ is defined by $\langle f_{0i} | \bar{s}s | 0 \rangle = m_{f_{0i}} \bar{f}_{f_{0i}}^s$.

It turns out that the nonresonant background in $B \rightarrow P_1 P_2$ transition described by Eq. (3) is too small to account for the experimental observation that $\overline{B}^0 \rightarrow K^+ K^- \overline{K}^0$ is dominated by the nonresonant contributions. This implies that the two-body matrix element e.g. $\langle K \overline{K} | \bar{s}s | 0 \rangle$ induced by the scalar density should have a large nonresonant component. In the absence of first-principles calculation, we will use the decay mode $\overline{B}^0 \rightarrow K_S K_S K_S$ or $B^- \rightarrow K^- K_S K_S$ to fix f_s^{NR} [14].

The calculated branching ratios of resonant and nonresonant contributions to $\overline{B}^0 \rightarrow K^+ K^- \overline{K}^0$ and $K_S \pi^0 \pi^0$ are summarized in Tables I and II, respectively. The theoretical errors shown there are from the uncertainties in (i) the parameter α_{NR} which governs the momentum dependence of the nonresonant amplitude [cf. Eq. (3)], (ii) the strange quark mass m_s , the form factor F_0^{BK} and the nonresonant contribution f_s^{NR} constrained by the $K_S K_S K_S$ rate, and (iii) the unitarity angle γ . We see that the predicted rates for resonant and nonresonant components are consistent with experiment. The nonresonant background arises dominantly from the transition process (90%) via the scalar density induced vacuum to $K \overline{K}$ transition, namely, $\langle K^+ K^- | \bar{s}s | 0 \rangle$, and slightly from the current-induced process (3%). In the $K_S \pi^0 \pi^0$ mode, the nonresonant fraction is found to be of order 40%. Recall that large nonresonant contributions to the decays $B^- \rightarrow K^- \pi^+ \pi^-$ and $\overline{B}^0 \rightarrow \overline{K}^0 \pi^+ \pi^-$, at the level of (35–40)%, have been reported by Belle [16].

The $K^+ K^- K_S$ mode is an admixture of CP -even and CP -odd components. By excluding the major CP -odd




FIG. 2: The K^+K^- mass spectra for $\bar{B}^0 \rightarrow K^+K^-K_S$ decay from (a) CP -even and (b) CP -odd contributions. The insert in (b) is for the ϕ region. The full $K^+K^-K_S$ spectrum, which is the sum of CP -even and CP -odd parts, measured by BaBar [9] is depicted in (c).

TABLE I: Branching ratios (in units of 10^{-6}) of resonant and nonresonant (NR) contributions to $\bar{B}^0 \rightarrow K^+K^-\bar{K}^0$. Theoretical errors correspond to the uncertainties in (i) α_{NR} , (ii) m_s , F_0^{BK} and σ_{NR} , and (iii) γ . Experimental results are taken from [9].

Decay mode	BaBar [9]	Theory
$\phi\bar{K}^0$	3.07 ± 0.45	$2.6^{+0.0+0.5+0.0}_{-0.0-0.4-0.0}$
$f_0(980)\bar{K}^0$	5.31 ± 2.19	$5.8^{+0.0+0.1+0.0}_{-0.0-0.5-0.0}$
$X_0(1550)K^-$	0.98 ± 0.44	$0.93^{+0.00+0.16+0.00}_{-0.00-0.15-0.00}$
NR	21.7 ± 5.1	$14.4^{+0.4+2.1+0.1}_{-0.5-1.5-0.1}$
total	$23.8 \pm 2.0 \pm 1.6$	$17.1^{+0.3+3.6+0.1}_{-0.3-2.9-0.1}$

TABLE II: Same as Table I except for the $\bar{K}^0\pi^0\pi^0$ mode.

Decay mode	Theory
$f_0(980)\bar{K}^0$	$3.8^{+0.0+2.0+0.0}_{-0.0-0.4-0.0}$
$\bar{K}^{*0}\pi^0$	$0.55^{+0.00+0.16+0.00}_{-0.00-0.13-0.00}$
$\bar{K}_0^{*0}(1430)\pi^0$	$2.3^{+0.0+0.8+0.0}_{-0.0-0.6-0.0}$
NR	$5.1^{+0.1+1.8+0.0}_{-0.1-1.0-0.0}$
total	$12.5^{+0.0+3.9+0.1}_{-0.0-2.9-0.1}$

contribution from ϕK_S , the 3-body $K^+K^-K_S$ final state is primarily CP -even. The K^+K^- mass spectra of the $\bar{B}^0 \rightarrow K^+K^-K_S$ decay from CP -even and CP -odd contributions are shown in Fig. 2. For the CP -even spectrum, there are peaks at the threshold and $m_{K^+K^-} = 1.5$ GeV region.¹ The threshold enhancement arises from the

TABLE III: Branching ratios for $\bar{B}^0 \rightarrow K^+K^-K_S$, $K_SK_SK_S$, $K_SK_SK_L$, $K_S\pi^0\pi^0$ decays and the fraction of CP -even contribution to $\bar{B}^0 \rightarrow K^+K^-K_S$. The branching ratio of CP -odd $K^+K^-K_S$ with ϕK_S excluded is shown in parentheses. Results for $(K^+K^-K_L)_{CP\pm}$ are identical to those for $(K^+K^-K_S)_{CP\mp}$. Experimental results are taken from [4].

Final State	$\mathcal{B}(10^{-6})_{\text{theory}}$	$\mathcal{B}(10^{-6})_{\text{expt}}$
$K^+K^-K_S$	$8.54^{+0.13+1.82+0.06}_{-0.14-1.45-0.06}$	12.4 ± 1.2
$(K^+K^-K_S)_{CP+}$	$6.97^{+0.04+1.37+0.04}_{-0.04-1.12-0.04}$	
$(K^+K^-K_S)_{CP-}$	$1.57^{+0.09+0.46+0.02}_{-0.10-0.32-0.02}$	
	$(0.14^{+0.06+0.14+0.01}_{-0.06-0.06-0.01})$	
$K_SK_SK_S$	input	6.2 ± 0.9
$K_SK_SK_L$	$6.06^{+0.11+0.61+0.02}_{-0.08-0.69-0.02}$	< 14
$K_S\pi^0\pi^0$	$6.24^{+0.01+1.96+0.03}_{-0.02-1.45-0.04}$	

$f_0(980)K_S$ and the nonresonant $f_S^{K^+K^-}$ contributions. For the CP -odd spectrum, the peak on the lower end corresponds to the ϕK_S contribution, which is also shown in the insert. The small middle hump in Fig. 2(a) comes from the interference between $b \rightarrow s$ and $b \rightarrow u$ amplitudes. The $b \rightarrow u$ transition is governed by the current-induced process $\langle \bar{B}^0 \rightarrow K^+\bar{K}^0 \rangle \times \langle 0 \rightarrow K^- \rangle$. From Eq. (3) it is clear that the $b \rightarrow u$ amplitude prefers a small invariant mass of K^+ and $\bar{K}^0$ and hence a large invariant mass of K^+ and K^- . In contrast, the $b \rightarrow c$ amplitude prefers a small s_{23} . The interference results to a minor hump shown in Fig. 2(a).

III. DECAY RATES AND CP ASYMMETRIES

Results for the decay rates and CP asymmetries in $\bar{B}^0 \rightarrow K^+K^-K_{S(L)}$, $K_SK_SK_{S(L)}$ are exhibited in Table III and Table IV, respectively. We see that

¹ In our previous work [17] we have argued that the spectrum should have a peak at the large $m_{K^+K^-}$ end. This is because we have introduced an additional nonresonant contribution to the ω_- parameter parametrized as $\omega_-^{NR} = \kappa \frac{2p_B \cdot p_2}{s_{12}^2}$ and employed the $B^- \rightarrow D^0 K^0 K^-$ data and applied isospin symmetry to the $\bar{B} \rightarrow K\bar{K}$ matrix elements to determine the unknown parameter κ . Since this nonresonant term favors a small $m_{K^+K_S}$ region, a peak of the spectrum at large $m_{K^+K^-}$ is thus expected. How-

ever, such a bump is not seen experimentally [9] [see also Fig. 2(c)]. In this work we will no longer consider this term.

TABLE IV: Mixing-induced and direct CP asymmetries $\sin 2\beta_{\text{eff}}$ (top) and A_f (in %, bottom), respectively, in $B^0 \rightarrow K^+ K^- K_S$, $K_S K_S K_S$ and $K_S \pi^0 \pi^0$ decays. Results for $(K^+ K^- K_L)_{CP\pm}$ are identical to those for $(K^+ K^- K_S)_{CP\mp}$. Experimental results are taken from [4].

Final state	$\sin 2\beta_{\text{eff}}$	Expt.
$(K^+ K^- K_S)_{\phi K_S}$ excluded	$0.721^{+0.000+0.001+0.009}_{-0.001-0.001-0.020}$	$0.58 \pm 0.13^{+0.12}_{-0.09}$
$(K^+ K^- K_S)_{CP+}$	$0.726^{+0.002+0.007+0.008}_{-0.002-0.004-0.019}$	
$(K^+ K^- K_L)_{\phi K_L}$ excluded	$0.721^{+0.000+0.001+0.009}_{-0.001-0.001-0.020}$	$0.58 \pm 0.13^{+0.12}_{-0.09}$
$K_S K_S K_S$	$0.719^{+0.000+0.000+0.008}_{-0.000-0.000-0.019}$	0.51 ± 0.21
$K_S K_S K_L$	$0.718^{+0.000+0.000+0.008}_{-0.000-0.000-0.019}$	
$K_S \pi^0 \pi^0$	$0.729^{+0.001+0.001+0.009}_{-0.001-0.001-0.019}$	$-0.84 \pm 0.71 \pm 0.08$
	$A_f(\%)$	Expt.
$(K^+ K^- K_S)_{\phi K_S}$ excluded	$-4.11^{+1.22+0.41+0.35}_{-0.91-0.42-0.30}$	-15 ± 9
$(K^+ K^- K_S)_{CP+}$	$-4.37^{+1.30+0.45+0.37}_{-1.00-0.46-0.31}$	
$(K^+ K^- K_L)_{\phi K_L}$ excluded	$-4.11^{+1.22+0.41+0.35}_{-0.91-0.42-0.30}$	-15 ± 9
$K_S K_S K_S$	$0.69^{+0.01+0.01+0.05}_{-0.01-0.01-0.06}$	23 ± 15
$K_S K_S K_L$	$0.90^{+0.00+0.00+0.06}_{-0.00-0.02-0.08}$	
$K_S \pi^0 \pi^0$	$0.32^{+0.05+0.02+0.02}_{-0.03-0.01-0.03}$	$-27 \pm 52 \pm 13$

the predicted branching ratios are consistent with the data within the theoretical and experimental errors. From Table IV we can calculate the deviation of the mixing-induced CP asymmetry in $B^0 \rightarrow K^+ K^- K_S$ and $K_S K_S K_S$ from that measured in $B \rightarrow \phi_{c\bar{c}} K_S$, namely, $\Delta \sin 2\beta_{\text{eff}} \equiv \sin 2\beta_{\text{eff}} - \sin 2\beta_{\phi_{c\bar{c}} K_S}$. Using the fitted CKM's $\sin 2\beta = 0.695^{+0.018}_{-0.016}$ [18] we obtain

$$\begin{aligned}
\Delta \sin 2\beta_{K^+ K^- K_S} &= 0.026^{+0.022}_{-0.030}, \\
\Delta \sin 2\beta_{K_S K_S K_S} &= 0.024^{+0.020}_{-0.025}, \\
\Delta \sin 2\beta_{K_S \pi^0 \pi^0} &= 0.034^{+0.020}_{-0.025}.
\end{aligned} \tag{7}$$

It should be stressed that despite the presence of color-allowed tree contributions in $B^0 \rightarrow K^+ K^- K_{S(L)}$, the deviation of the mixing-induced CP asymmetry in this penguin-dominated mode from that measured in $B \rightarrow \phi_{c\bar{c}} K_S$ is very similar to that of the $K_S K_S K_S$ mode. However, direct CP asymmetry of the former, being of order 4%, is more prominent than the latter.

Acknowledgments

I'm grateful to Chun-Khiang Chua and Amarjit Soni for fruitful collaboration.

-
- [1] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **71**, 091102 (2005).
[2] K. Abe *et al.* (Belle Collaboration), hep-ex/0507037.
[3] B. Aubert *et al.* (BaBar Collaboration), hep-ex/0507016.
[4] See web page of Heavy Flavor Averaging Group, <http://www.slac.stanford.edu/xorg/hfag>.
[5] D. London and A. Soni, Phys. Lett. B **407**, 61 (1997).
[6] Y. Grossman, Z. Ligeti, Y. Nir, and H. Quinn, Phys. Rev. D **68**, 015004 (2003).
[7] G. Engelhard, Y. Nir, and G. Raz, Phys. Rev. D **72**, 075013 (2005).
[8] M. Gronau and J.L. Rosner, Phys. Lett. B **564**, 90 (2003).
[9] B. Aubert *et al.* (BaBar Collaboration), hep-ex/0607112.
[10] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **74**, 032003 (2006).
[11] A. Garmash *et al.* (Belle Collaboration), Phys. Rev. D **71**, 092003 (2005).
[12] C. L. Y. Lee, M. Lu, and M. B. Wise, Phys. Rev. D **46**, 5040 (1992).
[13] H. Y. Cheng and K. C. Yang, Phys. Rev. D **66**, 054015 (2002).
[14] H.Y. Cheng, C.K. Chua, and A. Soni, in preparation.
[15] S.J. Brodsky and G.R. Farrar, Phys. Rev. D **11**, 1309 (1975).
[16] A. Garmash *et al.* (Belle Collaboration), Phys. Rev. Lett. **96**, 251803 (2006); Phys. Rev. D **75**, 012006 (2007).
[17] H.Y. Cheng, C.K. Chua, and A. Soni, Phys. Rev. D **72**, 094003 (2005).
[18] CKMfitter Group, J. Charles *et al.*, Eur. Phys. J. C **41**, 1 (2005) and updated results from <http://ckmfitter.in2p3.fr>; UTfit Collaboration, M. Bona *et al.*, hep-ph/0501199.